

Abstract Submitted  
for the 4CF15 Meeting of  
The American Physical Society

**Characterizing the Stellar Halo of M83**<sup>1</sup> MICHAEL BUSCH, Arizona State University, ERIC BELL, SARAH LOEBMAN, IAN ROEDERER, University of Michigan, GHOSTS TEAM<sup>2</sup> — Cosmological simulations of  $\Lambda$ CDM predict a hierarchical Galactic formation by the accretion of smaller satellite galaxies onto the main galactic plane. We observe far fewer satellite galaxies ( $\sim 20$ ) in the stellar halo in the Local Group than predicted by  $\Lambda$ CDM ( $\sim 500$ ). The GHOSTS (Galaxy Halos, Outer disks, Substructure, Thick disks and Star clusters) survey is the largest study of stellar populations in the stellar halos of 16 nearby disk galaxies using the Advanced Camera for Surveys (ACS) aboard the Hubble Space Telescope (HST). Additionally, the Subaru Telescope on Mauna Kea is a ground-based telescope capable of providing a wide-field survey of the stellar halo. This work characterizes the stellar halo of M83, a Milky Way type galaxy at a distance of  $\sim 5$  Mpc. We present early results on a method to use GHOSTS as a targeting survey to calibrate Subaru data, using Stellar Locus Regression to calculate color-color cuts and star-galaxy separation in Subaru data. The goal is to characterize the stellar populations, specifically Red Giant Branch (RGB) stars, as they are tracers of the underlying galactic formation.

<sup>1</sup>This work was supported by the National Science Foundation through the Research Experience for Undergraduates program. This research was completed in the Summer 2015 Physics REU at the University of Michigan.

<sup>2</sup>Galaxy Halos, Outer disks, Substructure, Thick disks and Star clusters

Michael Busch  
Arizona State University

Date submitted: 11 Sep 2015

Electronic form version 1.4