

Abstract Submitted
for the APR06 Meeting of
The American Physical Society

Spectrum of MHD Turbulence¹ STANISLAV BOLDYREV, University of Chicago — We propose a phenomenological theory of strong incompressible magnetohydrodynamic turbulence in the presence of a large-scale external magnetic field. We argue that in the inertial range of scales, magnetic-field and velocity-field fluctuations tend to align the directions of their polarizations. However, in driven turbulence the perfect alignment cannot be reached, it is precluded by the presence of a constant energy flux over scales. As a consequence, the directions of fluid and magnetic-field fluctuations at each scale λ become aligned within the angle $\phi_\lambda \propto \lambda^{1/4}$, which leads to scale-dependent depletion of nonlinear interaction and to the field-perpendicular energy spectrum $E(k_\perp) \propto k_\perp^{-3/2}$. Our results may be universal, i.e., independent of the external magnetic field, since small-scale fluctuations locally experience a strong field produced by large-scale eddies. The application of the results to diffractive interstellar scintillation is discussed.

¹Supported by the NSF Center for magnetic self-organization in Laboratory and Astrophysical Plasmas, U. Chicago; REF: astro-ph/0511290

Stanislav Boldyrev
University of Chicago

Date submitted: 12 Jan 2006

Electronic form version 1.4