

Abstract Submitted
for the APR06 Meeting of
The American Physical Society

Tidal Effects in the Resonance decay of light clusters¹ ROMUALDO DESOUZA, ALAN MCINTOSH, SYLVIE HUDAN, CARL METELKO, Dept. of Chemistry, Indiana University/IUCF — An excited nucleus can decay by emission of clusters. These clusters may be excited and themselves subsequently undergo particle decay. Peaks in the relative energy spectrum of the secondary decay products indicate resonance reflecting the discrete internal structure of the primary emitted cluster. Resonance spectroscopy can be used, for example, to determine the temperature of the initial source within a statistical approach. To date however, the effect of the field of the emitting nucleus on the decay of the cluster has been largely neglected. Tidal effects result in the correlation of the relative energy with emission angle as a function of the decay time. We explore the influence of the external Coulomb field on the decay of the first excited state of ^8Be in the reaction $^{60}\text{Ni} + ^{100}\text{Mo}$ at $E/A=11$ MeV. Experimental results will be compared with the predictions of a simple Coulomb trajectory model.

¹Support by the U.S. Department of Energy Grant No DE.FG02-88ER.40404

Romualdo deSouza
Dept. of Chemistry, Indiana University/IUCF

Date submitted: 12 Jan 2006

Electronic form version 1.4