

Abstract Submitted
for the APR06 Meeting of
The American Physical Society

A finite volume Fokker-Planck collision operator in constants-of-motion coordinates¹ Z. XIONG, X. Q. XU, B. I. COHEN, R. COHEN, M. R. DORR, J. A. HITTINGER, G. KERBEL, W. M. NEVINS, T. ROGNLIEN, LLNL — TEMPEST is a 5D gyrokinetic continuum code for edge plasmas. Constants of motion, namely, the total energy E and the magnetic moment μ , are chosen as coordinates because of their advantage in minimizing numerical diffusion in advection operators. Most existing collision operators are written in other coordinates; using them by interpolating is shown to be less satisfactory in maintaining overall numerical accuracy and conservation. Here we develop a Fokker-Planck collision operator directly in (E, μ) space using a finite volume approach. The (E, μ) grid is Cartesian, and the turning point boundary represents a straight line cutting through the grid that separates the physical and non-physical zones. The resulting cut-cells are treated by a cell-merging technique to ensure a complete particle conservation. A two dimensional fourth order reconstruction scheme is devised to achieve good numerical accuracy with modest number of grid points. The new collision operator will be benchmarked by numerical examples.

¹Work performed for USDOE by UC LLNL under contract W-7405-Eng-48

Z. Xiong
Lawrence Livermore National Laboratory

Date submitted: 13 Jan 2006

Electronic form version 1.4