

Abstract Submitted
for the APR06 Meeting of
The American Physical Society

Propagation of global shear Alfvén waves in gyrokinetic tokamak plasmas Y. NISHIMURA, Z. LIN, I. HOLOD, L. CHEN, UC-Irvine, V. DECYK, UCLA, S. KLASKY, ORNL, K. MA, UC-Davis, M. ADAMS, Columbia University, S. ETHIER, T. HAHM, W. LEE, J. LEWANDOWSKI, G. REWOLDT, W. WANG, PPPL — Employing the electromagnetic gyrokinetic simulation models,^{1,2} Alfvén wave dynamics in global tokamak geometry is studied. Based on a small parameter expansion by the square-root of the electron-ion mass ratio, the fluid-kinetic hybrid electron model² solves the adiabatic response in the lowest order and solves the kinetic response in the higher orders. We verify the propagation of shear Alfvén waves in the absence of drives or damping mechanisms by perturbing the magnetic field lines at $t = 0$ in a global eigenmode structure. The Alfvén wave experiences continuum damping.³ In the presence of energetic particles, excitations of toroidal Alfvén eigenmode (TAE) is expected within the frequency gap.⁴ With the η_i gradient drive, at a critical β value, the kinetic ballooning mode (KBM)⁵ is excited below the ideal MHD limit.

¹W.W.Lee *et al.*, Phys. Plasmas **8**, 4435 (2001).

²Z.Lin and L.Chen, Phys. Plasmas **8**, 1447 (2001).

³J.A.Tataronis and W. Grossman, Z. Phys. **14**, 203 (1973).

⁴C.Z.Cheng, L.Chen, and M.S.Chance, Ann.Phys. **161**, 21 (1984).

⁵C.Z.Cheng, Nucl. Fusion **22**, 773 (1982).

Yasutaro Nishimura
UC-Irvine

Date submitted: 13 Jan 2006

Electronic form version 1.4