

Abstract Submitted  
for the APR06 Meeting of  
The American Physical Society

**First Measurement of  $\sigma(gg \rightarrow t\bar{t})/\sigma(q\bar{q} \rightarrow t\bar{t})$  in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV<sup>1</sup>** SHABNAZ PASHAPOUR, Univ. of Toronto, CDF COLLABORATION  
— Following the discovery of the top quark, various studies have been dedicated to understanding the properties of this heavy particle. We will present the first measurement of the relative rate of  $t\bar{t}$  production through gluon-gluon fusion in quark-antiquark annihilation,  $\sigma(gg \rightarrow t\bar{t})/\sigma(q\bar{q} \rightarrow t\bar{t})$ . This measurement uses about  $0.7 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV collected by the Collider Detector at Fermilab (CDF). It provides a test of the QCD predictions for this pair-production mechanism, and a technique to test for unexpected sources of top quark production. The average number of gluons in the initial state of candidate  $t\bar{t}$  events is measured using the low  $p_T$  charged particle multiplicity, and the charged particle multiplicity is determined using W+n jet(s) and dijet data calibration samples, verified using Standard Model calculations.

<sup>1</sup>For the CDF Collaboration. This work has been supported by Natural Sciences and Research Council of Canada.

Robin Erbacher  
UC Davis

Date submitted: 16 Feb 2006

Electronic form version 1.4