

Abstract for an Invited Paper  
for the APR06 Meeting of  
The American Physical Society

### **Evolution of a Habitable Planet**

JAMES KASTING, Penn State University

Earth has remained habitable, and inhabited, over most of its 4.5-Gyr history despite an appreciable (30%) increase in solar luminosity over time and despite catastrophic events such as asteroid impacts and “Snowball Earth” episodes that have threatened biological survival on a global scale. Life has survived partly because of the resilience of the biota and partly because of feedback mechanisms that help to stabilize Earth’s global climate. In particular, buildup of volcanic CO<sub>2</sub> during times when the climate is cold provides a strong negative feedback that helps keep Earth within the temperature regime favorable to life. The same processes that help stabilize Earth’s climate should operate on other Earth-like planets, if they exist; thus, it is plausible that life could exist elsewhere. This hypothesis is now on the verge of being tested. NASA’s twin *Terrestrial Planet Finder (TPF)* missions, which could be launched as early as 2015-2020, will look for Earth-like planets around nearby stars and, if they are found, provide spectroscopic information on their atmospheres. Between them, these missions should be able to look for absorption bands of O<sub>2</sub>, H<sub>2</sub>O, CO<sub>2</sub>, and O<sub>3</sub>. Both O<sub>2</sub> and O<sub>3</sub> are considered to be good indicators of life for planets orbiting within the liquid water habitable zone of their parent star. NASA should be encouraged to give these missions high priority, so that we can answer these fundamental questions about the distribution of life in the universe.