

Abstract Submitted  
for the APR07 Meeting of  
The American Physical Society

**The  $\beta$  and delayed-neutron decay properties of neutron rich  $^{76-79}\text{Cu}$** <sup>1</sup> S.V. ILYUSHKIN, J.A. WINGER, Mississippi State, K.P. RYKACZEWSKI, C.J. GROSS, D. SHAPIRA, ORNL, R. GRZYWACZ, S.N. LIDDICK, C. MAZZOCCHI, S. PADGETT, M.M. RAJABALI, U. of Tennessee, J.C. BATCHELDER, UNIRIB/ORAU, A. KORGUL, Warsaw University, W. KROLÁS, Polish Academy of Sciences, E.F. ZGANJAR, A. PIECHAZEK, Louisiana State, C. GOODIN, J.H. HAMILTON, Vanderbilt University — Results of studies performed at the HRIBF on the  $\beta$ -decay of  $^{76-79}\text{Cu}$  isotopes are reported. Transmission through a charge exchange cell and isobar separator removed Zn from the ISOL beams and allowed fine tuning to obtain higher Cu concentrations. The  $\beta$ -delayed neutron probabilities of  $^{76}\text{Cu}$  and  $^{77}\text{Cu}$  were deduced from comparison of the intensities of  $\gamma$  rays in the Zn daughter isotopes and found to be 4(1)% and 38(4)%, respectively. For  $^{78}\text{Cu}$ , the yrast decay sequence transitions at 730, 891, and 907 keV suggest the highest feeding is to the  $6^+$  state with no population of the  $8^+$  isomer. For the r-process critical nuclide  $^{79}\text{Cu}$  we observed for the first time the 730 keV  $\gamma$  ray from the delayed-neutron branch thus solidifying this transition as the  $2^+ \rightarrow 0^+$  transition in  $^{78}\text{Zn}$ .

<sup>1</sup>This work was supported under US DOE grants DE-FG02-96ER41006, DE-AC05-00OR22725, DE-FG02-96ER40983, DE-AC05-06OR23100, DE-FG02-96ER40978, and DE-FG05-88ER40407.

Jeff Winger  
Mississippi State University

Date submitted: 16 Jan 2007

Electronic form version 1.4