

Abstract Submitted
for the APR09 Meeting of
The American Physical Society

Dynamic Confinement of ITER Plasma by O-Mode Driver at Electron Cyclotron Frequency Range V. ALEXANDER STEFAN, Nikola TESLA Laboratories (Stefan University), 1010 Pearl Street, La Jolla, CA 92038-2946. — A low B-field side launched electron cyclotron O-Mode driver¹ leads to the dynamic rf confinement, in addition to rf turbulent heating, of ITER² plasma. The scaling law for the local energy confinement time τ_E is evaluated ($\tau_E \sim 3n_e T_e / 2Q$, where $(3/2) n_e T_e$ is the local plasma thermal energy density and Q is the local rf turbulent heating rate). The dynamics of unstable dissipative trapped particle modes (DTPM)^{3,4} strongly coupled to Trivelpiece-Gould (T-G) modes is studied for gyrotron frequency 170GHz; power~24 MW CW; and on-axis B-field ~ 10 T. In the case of dynamic stabilization of DTPM turbulence and for the heavily damped T-G modes, the energy confinement time scales as $\tau_E \sim (I_0)^{-2}$, whereby I_0 (W/m²) is the O-Mode driver irradiance.

¹R. Prater et. al., *Nucl. Fusion* 48, No 3 (March 2008).

²E. P. Velikhov, History of the Russian Tokamak and the Tokamak Thermonuclear Fusion Research Worldwide That Led to ITER (Documentary movie; Stefan Studios Int'l, La Jolla, CA, 2008; © E. P. Velikhov, V. Stefan.)

³M N Rosenbluth, *Phys. Scr.* T2A 104-109 1982

⁴B. B. Kadomtsev and O. P. Pogutse, *Nucl. Fusion* 11, 67 (1971).

V. Alexander Stefan
Nikola TESLA Laboratories (Stefan University),
1010 Pearl Street, La Jolla, CA 92038-2946.

Date submitted: 12 Jan 2009

Electronic form version 1.4