

Abstract Submitted  
for the APR09 Meeting of  
The American Physical Society

**Magnetic Bubble Expansion as an Experimental Model for Extra-Galactic Radio Lobes** ALAN LYNN, YUE ZHANG, University of New Mexico, SCOTT HSU, HUI LI, WEI LIU, Los Alamos National Laboratory, MARK GILMORE, CHRISTOPHER WATTS, University of New Mexico — The Plasma Bubble Expansion Experiment (PBEX) has begun laboratory experiments and coordinated nonlinear MHD simulations to address outstanding nonlinear plasma physics issues related to how magnetic energy and helicity carried by extra-galactic jets interacts with the intergalactic medium to form radio lobe structures. Experiments are being conducted in the 4 meter long, 50 cm diameter HELCAT linear plasma device at UNM. A pulsed magnetized coaxial gun ( $\sim 10$  kV,  $\sim 100$  kA,  $\sim 2$  mWb) forms and injects magnetized plasma bubbles perpendicularly into a lower pressure weakly magnetized background plasma formed by a helicon and/or hot cathode source in HELCAT. Experimental parameters can be adjusted so that important dimensionless parameters are relevant to the astrophysical context. Ideal MHD simulations show that an MHD shock develops ahead of the bubble as it propagates, and that the bubble develops asymmetries due to the background field [1]. First experimental data from plasma bubble injection into a background plasma, including magnetic probe measurements and high-speed camera imaging, will be presented. [1] W. Liu et al., Phys. Plasmas 15, 072905 (2008). Supported by NSF-AST/DOE grant AST-0613577 and LANL LDRD.

Alan Lynn  
University of New Mexico

Date submitted: 09 Jan 2009

Electronic form version 1.4