

Abstract Submitted
for the APR09 Meeting of
The American Physical Society

Nonlinear Simulation of Alfvén Eigenmodes driven by Energetic Particles: Comparison between HMGC and TAEFL Codes¹ ANDREAS BIERWAGE, University of California, Irvine, DONALD A. SPONG, Oak Ridge National Laboratory, SCIDAC GSEP TEAM, ENEA/FRASCATI PLASMA THEORY TEAM — Hybrid-MHD-Gyrokinetic Code (HMGC) [1] and the gyrofluid code TAEFL [2,3] are used for nonlinear simulation of Alfvén Eigenmodes in Tokamak plasma. We compare results obtained in two cases: (I) a case designed for cross-code benchmark of TAE excitation; (II) a case based on a dedicated DIII-D shot #132707 where RSAE and TAE activity is observed. Differences between the numerical simulation results are discussed and future directions are outlined. [1] S. Briguglio, G. Vlad, F. Zonca and C. Kar, *Phys. Plasmas* 2 (1995) 3711. [2] D.A. Spong, B.A. Carreras and C.L. Hedrick, *Phys. Fluids B4* (1992) 3316. [3] D.A. Spong, B.A. Carreras and C.L. Hedrick, *Phys. Plasmas* 1 (1994) 1503.

¹This research is supported by U.S. DoE SciDAC GSEP.

Andreas Bierwage
University of California, Irvine

Date submitted: 12 Jan 2009

Electronic form version 1.4