

Abstract Submitted  
for the APR09 Meeting of  
The American Physical Society

**Reduction of the friction force in electron cooling systems due to magnetic field errors**<sup>1</sup> ANDREY SOBOL, GEORGE BELL, DAVID BRUH-WILER, Tech-X Corporation, ALEXEI FEDOTOV, VLADIMIR LITVINENKO, BNL — Magnetic field errors can limit the dynamical friction force on co-propagating ions and, hence, increase the cooling time. We present theoretical and numerical results for reduction of the friction force due to bounded transverse magnetic field errors, as a function of wavelength. VORPAL [1] simulations using a binary collision algorithm [2] show that small-wavelength field errors affect the friction logarithmically, via the Coulomb log, while long-wavelength errors reduce the friction by effectively increasing the transverse electron temperature. A complete understanding of finite-time effects and the role of small impact parameter collisions is required to correctly interpret the simulation results. We show that the distribution of electron-ion impact parameters is similar to a Pareto distribution, for which the central limit theorem does not apply. A new code has been developed to calculate the cumulative distribution function of electron-ion impact parameters and thus correctly estimate the expectation value and uncertainty of the friction force. [1] C. Nieter and J. Cary, *J. Comp. Phys.* 196 (2004), p. 448. [2] G. Bell et al., *J. Comp. Phys.* 227 (2008), p. 8714.

<sup>1</sup>Supported by the US DOE Office of Nuclear Physics under grants DE-FC02-07ER41499 and DE-FG02-04ER84094; used NERSC resources under grant DE-AC02-05CH11231

Andrey Sobol  
Tech-X Corporation

Date submitted: 13 Jan 2009

Electronic form version 1.4