

Abstract for an Invited Paper  
for the APR10 Meeting of  
The American Physical Society

**Highlights from Fermi Gamma-Ray Space Telescope observations of Gamma-Ray Bursts<sup>1</sup>**

JONATHAN GRANOT, University of Hertfordshire

The Fermi Gamma-Ray Space Telescope has more than doubled the number of Gamma-Ray Bursts (GRBs) detected at high energies ( $> 100$  MeV) within its first year of operation. Thanks to the very wide energy range covered by Fermi's Gamma-ray Burst Monitor (GBM; 8 keV to 40 MeV) and Large Area Telescope (LAT; 25 MeV to  $> 300$  GeV) it has measured the prompt GRB emission spectrum over an unprecedentedly large energy range (from  $\sim 8$  keV to  $\sim 30$  GeV). I will present highlights from Fermi GRB observations focusing mainly on the prompt emission phase. Interesting new observations will be discussed along with some of their possible implications, including: (i) What can we learn from the Fermi-LAT GRB detection rate, (ii) A limit on the variation of the speed of light with photon energy (for the first time beyond the Planck scale for a linear energy dependence from direct time of arrival measurements), (iii) Lower-limits on the bulk Lorentz factor of the GRB outflow (of  $\sim 1000$  for the brightest Fermi LAT GRBs), (iv) The detection (or in other cases, lack thereof) of a distinct spectral component at high (and sometimes also at low) energies, and possible implications for the prompt GRB emission mechanism, (v) The later onset (and longer duration) of the high-energy emission ( $> 100$  MeV) compared to the low-energy ( $< 1$  MeV) emission that is seen in most Fermi-LAT GRBs.

<sup>1</sup>J. Granot gratefully acknowledges a Royal Society Wolfson Research Merit Award.