

Abstract Submitted
for the APR10 Meeting of
The American Physical Society

Higgs Boson Exchange Effects in $l^+l^- \rightarrow W^+W^-$ at High Energy

ELENA BREWER, State University of New York - UB, ECC, Buffalo, NY 14261, USA, ULRICH BAUR, Department of Physics, State University of New York, Buffalo, NY 14260, USA — We consider the possibility for detecting effects due to the Higgs exchange diagram in high energy $\mu^+\mu^-$, e^+e^- , and $\tau^+\tau^-$ collisions producing a pair of W bosons. The processes $l^+l^- \rightarrow W^+W^-$ (with $l = \mu, e, \tau$) are analyzed, analytically and via numerical simulations, to determine the center of mass energy, $\sqrt{s_H}$, where the effects from Higgs exchange become relevant. The scaling of $\sqrt{s_H}$ with the mass of the incoming leptons is also studied. Special consideration is given to the $W^+W^- \rightarrow l^\pm\nu_l jj$ final state after experimental acceptance cuts are imposed. Angular cuts are shown to be able to significantly lower $\sqrt{s_H}$. The center of mass energy for which Higgs exchange effects become relevant in $l^+l^- \rightarrow W^+W^-$ is found to be much higher than that in $l^+l^- \rightarrow \bar{\nu}\nu W^+W^-$. Higgs boson exchange effects in W -pair production thus will not be relevant for the e^+e^- and $\mu^+\mu^-$ colliders currently on the drawing board.

Elena Brewer
State University of New York - UB, ECC, Buffalo, NY 14261

Date submitted: 22 Oct 2009

Electronic form version 1.4