

Abstract Submitted
for the APR10 Meeting of
The American Physical Society

D^0 Reconstruction in b -jets for Events with $Z^0 \rightarrow \mu^+\mu^- + b$ -jet
JESSICA METCALFE, I. GORELOV, S. SEIDEL, K. TOMS, University of New Mexico, ATLAS COLLABORATION — A Monte Carlo study on b -jet heavy quark fragmentation in ATLAS using fully reconstructed hadronic D mesons is presented. Events with a $Z^0 \rightarrow \mu^+\mu^-$ and a b -jet where $b \rightarrow D + X$ were studied. Several charmed hadronic modes were reconstructed. Presented here are the decay channels $D^0 \rightarrow K^-\pi^+$ and $D^{*+} \rightarrow D^0\pi^+, D^0 \rightarrow K^-\pi^+$. The mass resolution of the D^0 and D^{*+} is reported as well as the yield and reconstruction efficiencies for each species and the shape of the b -jet in terms of the p_T distribution of the D meson with respect to the jet axis. When these measurements are repeated with and compared to real LHC data, they will provide important information on the b -quark parton distribution inside the proton, and on the b -quark fragmentation distribution. Moreover, the analysis, when applied to real data, will give a better understanding of QCD backgrounds and provide an additional tool to constrain the b -jet energy scale.

Jaehoon Yu
Univ. of Texas at Arlington

Date submitted: 23 Oct 2009

Electronic form version 1.4