

Abstract Submitted  
for the APR12 Meeting of  
The American Physical Society

**Photoionization and Electron-Ion Recombination of Fe XVII for High Temperature Plasmas**<sup>1</sup> SULTANA NAHAR<sup>2</sup>, The Ohio State University — Earlier studies on electron-ion recombination of Fe XVII,  $e + Fe XVIII \rightarrow Fe XVII$ , concentrated on low temperature region. However, due to its higher abundance, recombination in the high temperature region such as one exists in solar convection zone is of great importance. Total and level-specific recombination cross sections and rates of Fe XVII are presented from the detailed study in the high temperature. The calculations were carried out using the unified method which incorporates both the radiative recombination (RR) and dielectronic recombination (DR) including the interference effects and provides self-consistent set of recombination rates and photoionization cross sections. The calculations included core excitations of levels n=2 and 3 complexes in contrast to earlier consideration of the n=2 levels only, and hence entails a large energy range and high temperature range. Resonances due to core excitations to n=3 levels are much more extensive and stronger than those to n=2 levels. These have introduced a broad and high FR peak at about  $5 \times 10^6$  K.

<sup>1</sup>Partially supported by DOE and NSF

<sup>2</sup>PDF presentation

Sultana Nahar  
The Ohio State University

Date submitted: 02 Feb 2012

Electronic form version 1.4