

Abstract Submitted
for the APR14 Meeting of
The American Physical Society

The thermonuclear reaction rate of $^{17}\text{O}(p,\gamma)^{18}\text{F}$ —a low-energy, high beam current study at LENA¹ MATTHEW BUCKNER, CHRISTIAN ILIADIS, KEEGAN KELLY, LORI DOWNEN, ARTHUR CHAMPAGNE, JOHN CESARATTO, RICHARD LONGLAND, The University of North Carolina at Chapel Hill, Triangle Universities Nuclear Laboratory, LENA TEAM — Classical novae are thought to be the dominant source of ^{17}O in our Galaxy. These energetic events produce ^{18}F that, as it decays to ^{18}O , drives the ejection of nuclear “ash” into the interstellar medium. The importance of the non-resonant component of the $^{17}\text{O}(p,\gamma)^{18}\text{F}$ reaction is well established, and numerous studies have been performed to analyze this reaction. However, the temperature regime relevant to explosive hydrogen burning during classical novae corresponds to very low proton bombarding energies. At these low energies, the Coulomb barrier suppresses the reaction yield in the laboratory, and environmental backgrounds dominate the detected signal making it difficult to differentiate the direct capture γ -cascade from background. At the Laboratory for Experimental Nuclear Astrophysics (LENA), our electron cyclotron resonance (ECR) ion source produces intense, low-energy protons (≈ 2.0 mA at the target), and these high currents boost the thermonuclear reaction yield. The LENA facility also has a coincidence detector setup that reduces environmental background contributions. Improved $^{17}\text{O}(p,\gamma)^{18}\text{F}$ direct capture reaction rates are currently being determined, and our progress will be reported.

¹The DOE NNSA Stewardship Science Graduate Fellowship under Grant no. DE-FC52-08NA28752.

Matthew Buckner
The University of North Carolina at Chapel Hill,
Triangle Universities Nuclear Laboratory

Date submitted: 10 Jan 2014

Electronic form version 1.4