

Abstract Submitted
for the APR15 Meeting of
The American Physical Society

Anisotropic hydrodynamics for conformal Gubser flow¹ MOHAMMAD NOPOUSH, Kent State Univ - Kent, RADOSLAW RYBLEWSKI², The H. Niewodniczanski Institute of Nuclear Physics, MICHAEL STRICKLAND, Kent State Univ - Kent — We derive the equations of motion for a system undergoing boost-invariant longitudinal and azimuthally-symmetric transverse “Gubser flow” using leading-order anisotropic hydrodynamics. This is accomplished by assuming that the one-particle distribution function is ellipsoidally-symmetric in the momenta conjugate to the de Sitter coordinates used to parameterize the Gubser flow. We then demonstrate that the $SO(3)_q$ symmetry in de Sitter space further constrains the anisotropy tensor to be of spheroidal form. The resulting system of two coupled ordinary differential equations for the de Sitter-space momentum scale and anisotropy parameter are solved numerically and compared to a recently obtained exact solution of the relaxation-time-approximation Boltzmann equation subject to the same flow. We show that anisotropic hydrodynamics describes the spatio-temporal evolution of the system better than all currently known dissipative hydrodynamics approaches. In addition, we prove that anisotropic hydrodynamics gives the exact solution of the relaxation-time approximation Boltzmann equation in the ideal, $\eta/s \rightarrow 0$, and free-streaming, $\eta/s \rightarrow \infty$, limits.

¹R.R. was supported by Polish National Science Center Grant No. DEC-2012/07/D/ST2/02125. M.S. was supported by U.S. DOE Award No. DE-AC0205CH11231

²This institute is a part of Polish Academy of Sciences, located at Krakow, Poland.

Mohammad Nopoush
Kent State Univ - Kent

Date submitted: 13 Dec 2014

Electronic form version 1.4