

Abstract Submitted
for the APR17 Meeting of
The American Physical Society

NOvA Short-Baseline Tau-Neutrino Appearance Search RIJEESH

KELOTH, Fermilab — Three-flavor neutrino oscillations have successfully explained a wide range of neutrino oscillation experiment results. However, anomalous results, such as the electron-antineutrino appearance excess seen by LSND and MiniBooNE, do not fit the three-flavor paradigm and can be explained by the addition of a sterile neutrino at a larger mass scale than the existing three flavor mass states. The NOvA experiment consists of two finely segmented, liquid scintillator detectors operating 14.6 mrad off-axis from the NuMI muon-neutrino beam. The Near Detector is located on the Fermilab campus, 1 km from the NuMI target, while the Far Detector is located at Ash River, MN, 810 km from the NuMI target. The NOvA experiment is primarily designed to measure electron-neutrino appearance at the Far Detector using the Near Detector to control systematic uncertainties; however, the Near Detector is well suited for searching for anomalous short-baseline oscillations. I will present a novel method for selecting tau neutrino interactions with high purity at the Near Detector using a convolutional neural network. Using this method, the sensitivity to anomalous short-baseline tau-neutrino appearance due to sterile neutrino oscillations will be presented.

Rijeesh Keloth
Fermilab

Date submitted: 29 Sep 2016

Electronic form version 1.4