

Abstract Submitted
for the APR17 Meeting of
The American Physical Society

Probing the CP nature of Higgs couplings in $t\bar{t}h$ events at the LHC MIGUEL FIOLHAIS, Borough of Manhattan Community College, CHRISTOPHER PEASE, City College of New York, City University of New York, ANTONIO ONOFRE, LIP, University of Minho — The CP nature of the Higgs coupling to top quarks ($t\bar{t}h$) is studied in proton-proton collision events at a centre-of-mass energy of 13 TeV at the LHC. Pure scalar and pseudo-scalar Higgs boson signal events are generated with MadGraph5_aMC@NLO, and analysed in dileptonic final states with two oppositely charged leptons and four jets. These events are fully reconstructed by applying a kinematic fit. As a result, new angular distributions of the decay products as well as CP angular asymmetries are explored to separate the scalar from the pseudo-scalar components of the Higgs boson, which allows to reduce the contribution from the dominant irreducible background, $t\bar{t}b\bar{b}$. In addition, significant differences between the angular distributions and asymmetries are observed, providing new observables for a global fit of the Higgs couplings parameters.

Miguel Fiolhais
Borough of Manhattan Community College, City University of New York

Date submitted: 30 Sep 2016

Electronic form version 1.4