

Abstract Submitted  
for the APR17 Meeting of  
The American Physical Society

**Chiral transport effects in charged anisotropic plasma within a magnetic field at strong coupling** MATTHIAS KAMINSKI, University of Alabama - Tuscaloosa, MARTIN AMMON, University of Jena, Germany, ROSHAN KOIRALA, University of Alabama - Tuscaloosa, JULIAN LEIBER, University of Jena, Germany, JACKSON WU, University of Alabama - Tuscaloosa — We compute the quasi-normal mode frequencies of gauge field and metric perturbations around black branes which are electrically and magnetically charged. By use of the gauge/gravity correspondence, these fluctuations are dual to conserved current operators of a particular class of strongly coupled field theories with a chiral anomaly. Within such a theory, we consider a thermal charged plasma state subjected to an external magnetic field. Quasi-normal mode frequencies are dual to the poles in the two-point functions of these conserved currents, encoding information about transport and dissipation in the plasma. For comparison, we also compute the same two-point functions in the hydrodynamic limit with field-theoretic methods. Together, these two approaches reveal various effects of the magnetic field and chiral transport coefficients on the location of the hydrodynamic poles (e.g. analogous to the chiral magnetic and chiral vortical effects), as well as transport effects beyond the hydrodynamic approximation. We conjecture qualitative conclusions for heavy-ion collision experiments.

Matthias Kaminski  
Univ of Alabama - Tuscaloosa

Date submitted: 30 Sep 2016

Electronic form version 1.4