

Abstract Submitted  
for the APR17 Meeting of  
The American Physical Society

**Extracting Neutron Polarizabilities from Compton Scattering on Quasi-Free Neutrons in  $\gamma d \rightarrow \gamma np$** <sup>1</sup> BERHAN DEMISSIE, George Washington Univ — Compton scattering processes are ideal to study electric and magnetic dipole polarizability coefficients of nucleons. These fundamental quantities parametrize the response to a monochromatic photon probe. In this work, the inelastic channel  $\gamma d \rightarrow \gamma np$  is treated in  $\chi$ EFT, with a focus on the NQFP - neutron quasi-free peak - kinematic region. In this region, the momentum of the outgoing proton is small enough that it is considered to remain at rest. This provides access to the Compton scattering process  $\gamma n \rightarrow \gamma n$  from which the neutron scalar polarizabilities  $\alpha$  and  $\beta$  are extracted. Using  $\chi$ EFT, differential cross-sections,  $d^3\sigma/dE_n d\Omega_{\gamma'} \Omega_n$ , in the photon energy range of 200-400 MeV are computed. The biggest contribution comes from the impulse approximation, with small corrections stemming from final state interaction, meson exchange currents and rescattering. A new extraction of neutron polarizabilities from a two-parameter fit to the Kossert et al. data taken at MAMI in 2002 is presented.

<sup>1</sup>This work is supported by the US Department of Energy under contracts DE-FG02-95ER-40907, and by the Deans Research Chair programme of the Columbian College of Arts and Sciences of The George Washington University.

Berhan Demissie  
George Washington Univ

Date submitted: 30 Sep 2016

Electronic form version 1.4