

Abstract Submitted  
for the APR21 Meeting of  
The American Physical Society

**Extracting twist three GPDs from deeply virtual compton scattering**<sup>1</sup> KYLE SHIELLS, Center for Nuclear Femtography, XIANGDONG JI, Center for Nuclear Femtography; University of Maryland, College Park, YUXUN GUO, University of Maryland, College Park — Observables in the hard exclusive lepton production of real photons can be cleanly expressed in terms of the compton amplitudes involving generalized parton distributions (GPDs). This process can be factorized into the product of a short-distance partonic subprocess with a long-distance, off forward hadronic matrix element. They involve nonlocal quark and gluon operators and are naturally expressed in terms of GPDs; quantities which may carry crucial information about the nucleon's intrinsic angular momentum. In this talk, I will be providing an overview of the theoretical formalisms of these processes and a detailed look at some of the relevant asymmetry observables is given to twist-three accuracy. In particular, the relevant twist-two and -three GPDs which complete a newly-found nucleon spin sum rule in the transverse polarization plane are discussed.

<sup>1</sup>This research is supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under contract number DE-SC0020682, and the Center for Nuclear Femtography, Southeastern Universities Research Association, Washington D.C.

YUXUN GUO  
University of Maryland, College Park

Date submitted: 07 Jan 2021

Electronic form version 1.4