

Abstract Submitted
for the APR21 Meeting of
The American Physical Society

High-energy neutrino and electromagnetic counterparts of post-merger jets from super massive black hole mergers¹ CHENGCHAO YUAN, KOHTA MURASE, Pennsylvania State University, SHIGEO KIMURA, Tohoku University, THEODORE B. ZHANG, PETER MESZAROS, Pennsylvania State University — Supermassive black hole (SMBH) coalescences are ubiquitous in the history of the Universe and often exhibit strong accretion activities and powerful jets. These SMBH mergers are also promising candidates for future gravitational wave detectors such as Laser Space Interferometric Antenna (LISA). In this work, we consider neutrino and electromagnetic counterpart emissions originating from the jet-induced shocks. We formulate the jet structures and relevant interactions therein, and then evaluate neutrino emission from each shock site. We find that month-to-year high-energy neutrino emission from the postmerger jet after the gravitational wave event is detectable by IceCube-Gen2 within approximately five to ten years of operation in optimistic cases where the cosmic-ray loading is sufficiently high and a mildly super-Eddington accretion is achieved. In addition, based on our model that predicts slowly fading transients with durations of $\sim 1 - 10$ months with a time delay from days to months after the coalescence, we discuss implications for EM follow-up observations after the GW detection.

¹C.C.Y. and P.M. acknowledge support from the Eberly Foundation. The work of K.M. is supported by the Alfred P. Sloan Foundation, NSF Grant No. AST-1908689, and KAKENHI No. 20H01901. B.T.Z. acknowledges the IGC fellowship. S.S.K. acknowledges the JSPS Research Fellowship, JSPS KAKENHI Grant No. 19J00198.

Chengchao Yuan
Pennsylvania State University

Date submitted: 08 Jan 2021

Electronic form version 1.4