

Abstract Submitted
for the APR21 Meeting of
The American Physical Society

Compton Polarimetry for the CREX Experiment¹ ALLISON ZEC,
Univ of Virginia, PREX-II/CREX COLLABORATION — The Jefferson Lab Continuous Electron Beam Accelerator Facility's experimental Hall A employs a Compton polarimeter to measure incoming beam polarization for parity violating electron scattering experiments. The polarimeter operates by amplifying green laser light in a Fabry-Perot cavity which then Compton scatters off the incoming electron beam. The scattered photons are then passed through a scintillating GSO (Gadolinium Oxyorthosilicate) crystal with a single photomultiplier tube. The polarization measurement is conducted by taking advantage of the helicity-dependence of Compton scattering. By measuring the integrated signal from photons scattered while the beam is in different helicity states, we generate a differential asymmetry between these states, which then yields information about the electron beam's longitudinal polarization. Measuring the asymmetry requires a robust background subtraction of helicity-correlated asymmetry as well as identifying the Compton edge from observing spectra. The beam polarization is an important component needed to determine the parity-violating asymmetry for the CREX experiment. This talk will focus on the analysis of the integrating photon detector data.

¹NSF/DOE Grant

Allison Zec
Univ of Virginia

Date submitted: 23 Mar 2021

Electronic form version 1.4