

Abstract Submitted
for the DAMOP06 Meeting of
The American Physical Society

High harmonic generation from ions in a capillary discharge¹

DAVID M. GAUDIOSI, BRENDAN REAGAN, TENIO POPMINTCHEV, MICHAEL GRISHAM, MARK BERRILL, OREN COHEN, BARRY C. WALKER, University of Delaware, MARGARET M. MURNANE, HENRY C. KAPTEYN, University of Colorado, JORGE J. ROCCA, Colorado State University, NSF ERC FOR EXTREME ULTRAVIOLET SCIENCE AND TECHNOLOGY COLLABORATION — We demonstrate a significant extension of the high harmonic cutoff observed in xenon, up to 150 eV, by generating harmonics from ions in a capillary discharge plasma. Usually, the highest harmonics that can be generated from an atom or ion are limited by ionization-induced defocusing, beam attenuation of the laser, and a lack of phase matching of the frequency conversion process. Using a pre-ionized plasma generated by the capillary discharge, we dramatically reduce ionization-induced defocusing and energy loss of the driving laser due to ionization, allowing higher photon energies to be generated from xenon ions. We also demonstrate enhancement of the harmonic flux of nearly two orders of magnitude at photon energies around 90 eV when the capillary discharge is used to ionize xenon, compared with harmonic generation in a hollow waveguide. The use of a capillary discharge plasma for high harmonic generation shows great promise for extending efficient harmonic generation to much shorter wavelengths.

¹US Dept. of Energy, Office of Basic Energy Science, and the Engineering Research Centers Program of the National Science Foundation

David M. Gaudiosi
JILA/University of Colorado

Date submitted: 06 Feb 2006

Electronic form version 1.4