

Abstract Submitted
for the DAMOP07 Meeting of
The American Physical Society

Theory of x-ray absorption by laser-dressed atoms¹ CHRISTIAN BUTH, ROBIN SANTRA, Argonne National Laboratory, Argonne, Illinois 60439, USA — We present an *ab initio* theory for the x-ray photoabsorption cross section of atoms in the field of a moderately intense optical laser (800 nm, $10^{13} \frac{\text{W}}{\text{cm}^2}$). The laser dresses the core-excited atomic states, which introduces a dependence of the cross section on the angle between the polarization vectors of the two linearly polarized radiation sources. The strong interaction due to the laser-dressing is treated by diagonalization of a Floquet-type matrix; the weak coupling between x-rays and the atom is described by non-Hermitian perturbation theory. We apply our theory to study the photoabsorption cross section of neon and krypton atoms near the K edge. A pronounced modification of the cross section is found in the presence of the optical laser — **reference:** arXiv:physics/0611122.

¹C.B. is self-employed (Germany) and was funded by a Feodor Lynen Research Fellowship from the Alexander von Humboldt Foundation. R.S.'s work was supported by the Office of Science, U.S. Department of Energy, under Contract No. DE-AC02-06CH11357

Christian Buth

Date submitted: 30 Jan 2007

Electronic form version 1.4