

Abstract Submitted
for the DAMOP07 Meeting of
The American Physical Society

“Step-up” vs. “Step-sown” scattering asymmetry in the charge transfer of H^- on free-electron vicinal metallic surfaces BOYAN OBRESHKOV, UWE THUMM, Dept. of Physics, Kansas State University — We present numerical results based on a wave-packet propagation study of the one-electron charge transfer between H^- ions and free-electron vicinal metallic surfaces [1]. We derive an effective potential for the motion of the active electron within a Thomas-Fermi-von Weizsäcker model and extend this model to include the image charge effects. We first calculate H^- affinity level shift and width in fixed-ion approximation and solve a rate equation for the ion-survival probability for projectiles that are incident with a kinetic energy of 50 eV. We find an enhancement of the electron loss near the steps of the surface, due to the Smoluchowski effect. As a consequence, the ion-survival is more likely if the projectiles approach steps from above than from below [2].

[1] B. Obreshkov and U. Thumm, Phys. Rev. A **74**, 012901 (2006).

[2] B. Obreshkov and U. Thumm, Surf. Sci. **601**, 622 (2007).

Supported by NSF and the Division of Chemical Sciences, Office of Basis Energy Sciences, Office of Energy Research, US DoE.

Boyan Obreshkov
Dept. of Physics, Kansas State University

Date submitted: 02 Feb 2007

Electronic form version 1.4