

Abstract Submitted
for the DAMOP08 Meeting of
The American Physical Society

Revisited LeRoy-Bernstein formula for weakly bound molecules

LAURENCE PRUVOST, HAIKEL JELASSI, BRUNO VIARIS DE LESEGNO, lab Aimé Cotton, cnrs Orsay, LAC TEAM — The energy law giving the eigen energies of a $-cn/Rn-cm/Rm$ potential (studied by LeRoy in 1980 [1]) is revisited. For $n=3$, $m=6$, an analytical law giving the density of states is deduced. In the context of weakly bound levels an energy law giving the vibrational number v versus the binding energy is given. We show that the well-known LeRoy-Bernstein formula [2] [3] has to be corrected by additional terms, with the first one varying as the binding energy, the second one as its square and the third as a power to $7/6$. The use of such a law is discussed in the context of the photo-association spectroscopy of long range molecular levels. References [1] Theory of deviations from the limiting near-dissociation behaviour of diatomic molecules, R. J. LeRoy, J. Chem. Phys. 73, 6003 (1980). [2] Dissociation energy and long-range potential of diatomic molecules from vibrational spacings of higher levels, R. J. LeRoy and R. B. Bernstein, J. Chem. Phys. 52, 3869 (1970). [3] The Dissociation Energy of the Hydrogen Molecule Using Long-Range Forces, W. C. Stwalley, Chem. Phys. Lett. 6, 241,1970.

Laurence Pruvost
lab Aimé Cotton, cnrs Orsay

Date submitted: 28 Jan 2008

Electronic form version 1.4