

Abstract Submitted  
for the DAMOP08 Meeting of  
The American Physical Society

**Valence calculations of binding energies of mid-row lanthanides<sup>1</sup>**

STEVEN M. O'MALLEY, DONALD R. BECK, Physics Department, Michigan Technological University — Binding energies of mid-row lanthanides have been calculated by taking advantage of the >90%  $LS$  purity of the  $4f^n$  subgroups of their neutral ground states <sup>2</sup>. As described in our recent work on  $\text{Nd}^{-3}$ , the  $4f^n$  electrons were treated as core-like and restricted to a single  $LS$  term throughout all Dirac-Fock and correlation configurations, resulting in manageable relativistic configuration-interaction calculations for these computationally complex systems. Several weakly bound  $6p$  attachments (<200 meV) were found for each negative ion, and these were carefully analyzed with respect to approximate  $LS$  total configuration as well as the  $j$ 's of the  $4f^n$  subgroup and  $6p$  electron.

<sup>1</sup>Supported by the National Science Foundation, Grant No. PHY-0097111.

<sup>2</sup>*Atomic Energy Levels – The Rare-Earth Elements*, edited by W. C. Martin, R. Zalubas, and L. Hagan, Natl. Bur. Stand. Ref. Data Ser. Natl. Bur. Stand. (U.S.) Circ. No. 60 (U.S. GPO, Washington, D.C., 1978).

<sup>3</sup>S. M. O'Malley and D. R. Beck, *Phy. Rev. A* 77, 012505 (2008).

Steven M. O'Malley  
Physics Department, Michigan Technological University

Date submitted: 04 Feb 2008

Electronic form version 1.4