

Abstract Submitted  
for the DAMOP08 Meeting of  
The American Physical Society

**Highly Directional Emission from Inhomogeneous Dielectric Microcavities** G. PAINCHAUD-APRIL, J. POIRIER, Uni. Laval (Québec), S. SAIDI, Y.-A. PETER, Ecole Polytechnique (Montréal), E. BRASSELET, Uni. Bordeaux I (France), L.J. DUBÉ, Uni. Laval (Québec) — We propose a novel method of extracting light beams from 2D microcavities. The concept is based on *inhomogeneous dielectric cavities* (IDC) where the inhomogeneities arise from a space-dependent refractive index whose variations may be continuous (e.g. a localized induced gaussian profile of the index) or discontinuous (e.g. holes or refractive steps in the cavity material). Instead of smooth deformations of a circular cavity that produce directional output while sacrificing the quality factor  $Q$ , we intend to operate with an integrable geometry (a disk) and induce directionality through the (possibly reconfigurable) medium while preserving a high  $Q$ . The systems are interesting on two counts. Firstly, as classical objects, the IDC are equivalent to dielectric billiards (i.e. photonic escape is possible) where the broken symmetry of the material can induce a transition from regular to chaotic dynamics: chaos in an integrable billiard geometry. Secondly, guided by the classical phase space information, the wave dynamics can be “engineered” to produce highly directional emission with tailored optical properties, the grail of microcavity research. We have studied a number of configurations and will present results on their respective performances.

Louis J. Dubé  
Université Laval

Date submitted: 04 Feb 2008

Electronic form version 1.4