

Abstract Submitted  
for the DAMOP09 Meeting of  
The American Physical Society

**Atomic mass measurements for neutrino mass**<sup>1</sup> MATTHEW REDSHAW, BRIANNA MOUNT, EDMUND MYERS, Florida State University — As usually understood, observation of neutrinoless double-beta-decay implies that neutrinos are their own antiparticles (Majorana particles), while measurements of the decay rate, or limits on the rate, provide information on absolute neutrino mass. Large-scale neutrinoless double-beta-decay detectors, proposed or under development, such as EXO, CUORE, GERDA, MAJORANA, etc. should be sensitive to a linear combination of neutrino masses, the “effective Majorana mass of the electron neutrino”, below 0.1 eV/ $c^2$ . The signature of neutrinoless double-beta decay is a sharp peak in the total electron-energy spectrum at the Q-value of the decay – the mass-energy difference between the parent and daughter atoms. Using one or two multiply-charged ions in a Penning trap, we have now measured the atomic masses of  $^{136}\text{Xe}$ ,  $^{130}\text{Te}$ ,  $^{130}\text{Xe}$ ,  $^{76}\text{Ge}$ ,  $^{76}\text{Se}$  to a fractional precision of  $2 \times 10^{-10}$  or better, corresponding to Q-values with uncertainties below 25 eV. This is more than sufficient precision for the proposed large-scale experiments. Progress on mass measurements of  $^{74}\text{Ge}$  and  $^{74}\text{Se}$ , relevant to resonance-enhanced neutrinoless double-electron capture in  $^{74}\text{Se}$ , will also be reported.

<sup>1</sup>Support from NSF and NIST

Edmund Myers  
Florida State University

Date submitted: 22 Jan 2009

Electronic form version 1.4