

Abstract Submitted
for the DAMOP09 Meeting of
The American Physical Society

Using hyperfine structure to investigate perturbations between highly-excited states: the HF C-X spectrum¹ JEFFREY PHILIPPSON, RALPH SHIELL, Trent University, ELMAR REINHOLD, WIM UBACHS, Vrije Universiteit, Amsterdam, NL — It has long been known that the B $^1\Sigma^+$ ion-pair state in HF is strongly perturbed by electronic Rydberg states [1]. We present a quantitative analysis of these perturbations through their effects on the fluorine orbital magnetic hyperfine parameter obtained from XUV spectra of the C $^1\Pi$, $v=0$ -X $^1\Sigma^+$, $v=0$ transition [2]. A Λ -doubling interaction between the ground vibrational level of the C-state and the nearby $v=29$ level of the B-state produces an apparent rotational state dependence in the values of this parameter derived from the R-branch lines. This work demonstrates how insight into the extent of inter-state perturbations can be obtained from the variation of hyperfine parameters.

[1] A. E. Douglas and F. R. Greening, *Can. J. Phys.* **57**, 1650 (1979).

[2] J. N. Philippson, R. C. Shiell, E. Reinhold and W. Ubachs, *J. Chem. Phys.* **129**, 174310 (2008).

¹Research supported by the European Community Integrated Infrastructure Initiative action LASERLAB-Europe and NSERC (Canada).

Jeffrey Philippson
Trent University

Date submitted: 26 Jan 2009

Electronic form version 1.4