

Abstract Submitted
for the DAMOP09 Meeting of
The American Physical Society

SEOP polarization of ^3He and ^{129}Xe at high volumetric rates¹

I. RUSET², Xemed LLC and University of New Hampshire, D. WATT, J. DISTELBRINK, J. BRACKETT, J. KETEL, S. KETEL, W. PORTER, Xemed LLC, Durham, New Hampshire, T. UPTON, University of New Hampshire, W. HERSMAN³, University of New Hampshire and Xemed LLC — New regimes for production of polarized ^3He and ^{129}Xe using Spin Exchange Optical Pumping are under study for the purpose of increasing volumetric production rates, output polarization, or both. For polarized ^{129}Xe production, photon efficiency is increased by polarizing at low xenon partial pressure. Higher spin exchange rates between alkali atoms and the xenon nucleus are accessed by exploiting longer lived van der Waals molecules at lower buffer gas pressures. High production rates are accomplished by flowing at high volumetric rates and illuminating with a high power, narrowed line laser. Polarizations of ^{129}Xe exceeding 70% at production rates of 3 SLPH were observed. For polarized ^3He production, efficiency is maximized by selecting a high operating temperature and an alkali mixture rich in potassium. High volumetric production rate is accomplished by illuminating a large 8 L high-pressure ^3He reservoir with a high power (broad) diode stack. We estimate that 50 L can be polarized through three exponential ramp-up times in 12 hours, yielding as much as 100 L/day. Asymptotic ^3He polarization will be reported.

¹Supported by NIH grants R44HL87550, R44EB07439; and DOE grants FG02-88ER40410, FG02-08ER86368 and FG02-08ER86369.

²Xemed is primary affiliation

³UNH is my primary institution

William Hersman
University of New Hampshire and Xemed LLC

Date submitted: 26 Jan 2009

Electronic form version 1.4