

Abstract Submitted
for the DAMOP09 Meeting of
The American Physical Society

Theoretical Studies of Dissociative Recombination¹ D.O. KASHINSKI, R.F. MALEND, A.P. HICKMAN, Lehigh University, D. TALBI, Université Montpellier II — We are currently investigating the dissociative recombination (DR) of electrons with the molecular ions N_2H^+ and C_3H_3^+ . These ions exist in the interstellar medium, and accurate DR rate constants are needed for astrophysical models. We are performing large scale electronic structure calculations of the excited-state potential surfaces of N_2H necessary to treat the process $e^- + \text{N}_2\text{H}^+ \rightarrow \text{N}_2 + \text{H}$ or $\text{N} + \text{NH}$. The work is based on using the block diagonalization method to determine diabatic potential curves. The dissociating curve that governs DR is then easily identified, and off-diagonal coupling terms can be used to determine the autoionization width Γ that is essential for a dynamics calculation. The status of the calculations will be presented at the conference. We have also investigated the normal modes of the molecular ion C_3H_3^+ . We expect that energy flow into and out of the vibration of a single CH bond may influence the overall DR dynamics, and we account for this effect using an appropriate quantum mechanical wave function for the initial state.

¹Work supported by NSF and the TeraGrid.

David O. Kashinski
Lehigh University

Date submitted: 26 Jan 2009

Electronic form version 1.4