

Abstract Submitted
for the DAMOP09 Meeting of
The American Physical Society

An Electron Electric Dipole Moment Search in the $^3\Delta_1$ Ground State of Tungsten Carbide Molecules JEONGWON LEE, AARON LEANHARDT, Department of Physics, University of Michigan, Ann Arbor, Michigan 48109-1040, USA — We report on progress towards constructing and characterizing a continuous tungsten carbide (WC) molecular beam for an electron electric dipole moment (EDM) search¹. Tungsten atoms are evaporated from a resistively heated filament and tungsten carbide molecules are formed through a reaction with methane: $W + CH_4 \rightarrow WC + 2H_2$. WC has a $^3\Delta_1$ ground state with its two valence electrons in a $\sigma\delta$ molecular orbital configuration^{2,3,4}. This molecular structure has several unique advantages for an electron EDM search and arises in other diatomic species such as $HfF^{+5,6}$, $ThF^{+5,7}$, and $ThO^{7,8}$.

¹J. Lee et al., Atomic Physics **XXI**, 190 (2008).

²K. Balasubramanian, J. Chem.Phys.**112**, 7425 (2000).

³S. M. Sickafoose et al., J. Chem. Phys. **116**, 993 (2002).

⁴D. Rothgeb et al., J. Chem.Phys.**129**, 114304 (2008).

⁵<http://jilawww.colorado.edu/bec/CornellGroup/>.

⁶E.R.Meyer et al., Phys. Rev. A **73**, 062108 (2006).

⁷E. R. Meyer et al., Phys. Rev. A **78**, 010502 (2008).

⁸A. C.Vutha et al., Atomic Physics **XXI**, 191 (2008).

Aaron Leanhardt
Department of Physics, University of Michigan,
Ann Arbor, Michigan 48109-1040, USA

Date submitted: 27 Jan 2009

Electronic form version 1.4