

Abstract Submitted
for the DAMOP10 Meeting of
The American Physical Society

Imaging of spatial correlations of Rydberg excitations in cold atom clouds¹ ANDREW SCHWARZKOPF, RACHEL SAPIRO, GEORG RAITHEL, University of Michigan — Previously, Rydberg excitation blockades have been shown to cause a saturation of Rydberg excitation numbers in atom samples and a narrowing of the excitation number statistics, and they have been employed in quantum information experiments. In the experiment described in this poster, it is planned to measure structures in the Rydberg pair correlation function predicted in.² To achieve sufficient spatial magnification, we use the principle of field ion microscopy. A tungsten tip is placed close to a cold atom cloud in which several Rydberg excitations are prepared using a narrow-linewidth laser. To read out the sample, the tip voltage is suddenly switched to a high value. The Rydberg atoms are field-ionized, and the resultant ions are projected onto a nearby position-sensitive detector. All components of this experiment are operational, as will be shown on this poster. Progress towards the imaging of spatial correlations of excitations will be discussed.

¹We gratefully acknowledge support from AFOSR and NSF-FOCUS.

²F. Robicheaux and J. Hernandez, “Many-body wave function in a dipole blockade configuration,” Phys. Rev. **A 72**, 63403, 1-4 (2005).

Andrew Schwarzkopf
University of Michigan

Date submitted: 27 Jan 2010

Electronic form version 1.4