

DAMOP12-2012-000043

Abstract for an Invited Paper
for the DAMOP12 Meeting of
the American Physical Society

Dipolar Chromium BECs

BRUNO LABURTHER-TOLRA, CNRS

Bose-Einstein condensates (BECs) made of ^{52}Cr atoms reveal new phenomena, due to the presence of the long-range and anisotropic dipole-dipole interactions (see for example [1]). In this talk, I will describe the effect of dipolar interactions on the properties of multi-component (spinor) Cr condensates at extremely low magnetic fields. Due to its anisotropy, the dipolar interaction introduces magnetization-changing collisions, which dynamically freeze the magnetization of the gas. We have thus observed a demagnetization of the BEC when the magnetic field is quenched below a critical value B_c corresponding to a phase transition between a ferromagnetic and a non-polarized ground state. The phase transition is due to an inter-play between spin-dependent interactions and the linear Zeeman effect [2]. We have also studied the thermodynamic properties of spinor Cr atoms, and we have observed that above the critical field B_c , the ferromagnetic nature of BECs leads to the spontaneous magnetization of the cloud when BEC is reached [3]. I will also describe the control of magnetization-changing collisions in optical lattices. We investigate a scheme in which dipolar relaxation is resonant when the energy released in dipolar relaxation matches a band excitation resonance [4]. This scheme, which may produce correlated pairs of rotating states in each lattice site, can be viewed as the equivalent of the Einstein-de-Haas effect. Although rotation is not yet produced in our experiment, I will present first experimental results of these dipolar resonances, which show a pronounced anisotropic behaviour.

[1] T. Lahaye et al., Rep. Prog. Phys. 72, 126401 (2009), G. Bismut, et al., Phys. Rev. Lett. 105, 040404 (2010)

[2] B. Pasquiou et al., Phys. Rev. Lett. 106, 255303 (2011)

[3] B. Pasquiou, arXiv:1110.0786, to be published in Phys. Rev. Lett. (2012)

[4] B. Pasquiou et al., Phys. Rev. Lett. 106, 015301 (2011)