

Abstract Submitted
for the DAMOP12 Meeting of
The American Physical Society

Trapped atom number in millimeter-scale magneto-optical traps

GREGORY W. HOTH, ELIZABETH A. DONLEY, JOHN KITCHING, National Institute of Standards and Technology — For compact cold-atom instruments, it is desirable to trap a large number of atoms in a small volume to maximize the signal-to-noise ratio. In MOTs with beam diameters of a centimeter or larger, the slowing force is roughly constant versus velocity and the trapped atom number scales as d^4 . For millimeter-scale MOTs formed from pyramidal reflectors, a d^6 dependence has been observed [Pollack et al., Opt. Express **17**, 14109 (2009)]. A d^6 scaling is expected for small MOTs, where the slowing force is proportional to the atom velocity. For a 1 mm diameter MOT, a d^6 scaling results in 10 atoms, and the difference between a d^4 and a d^6 dependence corresponds to a factor of 1000 in atom number and a factor of 30 in the signal-to-noise ratio. We have observed $> 10^4$ atoms in 1 mm diameter MOTs, consistent with a d^4 dependence. We are currently performing measurements for sub-mm MOTs to determine where the d^4 to d^6 crossover occurs in our system. We are also exploring MOTs based on linear polarization, which can potentially produce stronger slowing forces due to stimulated emission [Emile et al., Europhys. Lett. **20**, 687 (1992)]. It may be possible to trap more atoms in small volumes with this method, since high intensities can be easily achieved.

Elizabeth Donley
National Institute of Standards and Technology

Date submitted: 27 Jan 2012

Electronic form version 1.4