

Abstract Submitted
for the DAMOP12 Meeting of
The American Physical Society

Mechanisms of dissociative recombination of N_2H^+ and HCO^+ ¹

SAMANTHA FONSECA DOS SANTOS, NICOLAS DOUGUET, ANN OREL, UC-Davis, VIATCHESLAV KOKOULINE, UCF — N_2H^+ and HCO^+ are among the first molecular ions observed in the interstellar medium. They are both formed by fast proton transfer mechanisms and are destroyed either by taking part on the molecular synthesis of more complex molecules or by dissociative recombination (DR). Given the astrophysical importance of these ions, their DR rates have been the subject of several different studies over the years. Recently, good agreement between theory and experiment have been achieved for HCO^+ , however this is not the case for N_2H^+ . This is mainly due to the fact that previous theories only considered the direct DR mechanism in their calculations. Applying a general simplified model based on multichannel quantum defect theory that accounts for all the main ingredients of indirect DR we calculate cross sections and DR rates for N_2H^+ as well as for HCO^+ using an improved coupling scheme.

¹This work was supported the DOE, Office of Basic Energy Science and the National Science Foundation, Grant No's. PHY-08-55092 and PHY-08-55622

Samantha Fonseca dos Santos
UCDavis

Date submitted: 27 Jan 2012

Electronic form version 1.4