

Abstract Submitted  
for the DAMOP12 Meeting of  
The American Physical Society

**Few-photon optical nonlinearities using cold Rydberg atoms**

THIBAULT PEYRONEL, MIT, OFER FIRSTENBERG, Harvard University, QIYU LIANG, VLADAN VULETIC, MIT — Effects of the Rydberg blockade in cold atomic clouds have been intensively explored over the last few years. Optical fields can be coherently mapped onto atomic states with a Rydberg component using EIT techniques. As the dipole-dipole interaction between Rydberg atoms prevents several polaritons from propagating simultaneously within a Rydberg volume, it gives rise to strong non-linearities which are mapped back on the probe optical field. We aim at bringing the Rydberg-EIT nonlinear features into the single-photon regime in order to produce non-classical highly correlated states of light. Rubidium atoms are loaded in a far off-resonant (1064nm) optical dipole trap, where densities are typically large enough to reach high optical depths within a single blockade volume. In this regime, the outcoming photon-photon correlation function is expected to exhibit highly non-classical behavior, corresponding to trains of spatially separated single-photons. Moreover, EIT techniques together with a high-resolution imaging system allow the observation of Rydberg excitations in the quasi-1D configuration, and should pave the way to in-situ monitoring of strongly correlated many-body states such as the crystallisation of Rydberg atoms.

Thibault Peyronel  
MIT

Date submitted: 01 Feb 2012

Electronic form version 1.4