

Abstract Submitted
for the DAMOP13 Meeting of
The American Physical Society

Comparison of positron and electron binding to molecules¹ J.R. DANIELSON, A.C.L. JONES, M.R. NATISIN, C.M. SURKO, University of California, San Diego — Positrons can attach to molecules via Feshbach resonances in which a vibrational mode absorbs the excess energy. Using a high-resolution positron beam, this process has been used to measure positron-molecule binding energies for many chemical species.² Recent measurements have focused on molecules with large permanent dipole moments (i.e., $\mu > 2.5$ D), including aldehydes, ketones, and nitriles.³ Positron binding to these molecules is compared to the analogous weakly bound electron-molecule states, commonly referred to as “dipole-bound” negative anions.⁴ Positron binding energies are found to be one to two orders of magnitude larger than those of the analogous negative ions due to two effects: the orientation of the molecular dipole allows the positron to approach it more closely; and for positrons, lepton correlations (e.g., via polarizability) contribute more strongly.⁵ Comparisons to available calculations will be presented, as will comparisons to binding to molecules with $\mu \sim 0$ (e.g., polarizability bound states).

¹Work supported by NSF grant PHY 10-68023.

²G. F. Gribakin, et al., RMP **82**, 2557 (2010).

³J. R. Danielson, et al., PRA **85**, 022709 (2012).

⁴N. I. Hammer, et al., JCP **119**, 3650 (2003).

⁵J. R. Danielson, et. al, PRL **109**, 113201 (2012)

James Danielson
University of California, San Diego

Date submitted: 22 Jan 2013

Electronic form version 1.4