

Abstract Submitted  
for the DAMOP13 Meeting of  
The American Physical Society

**Ultrafast entanglement of trapped ions**<sup>1</sup> BRIAN NEYENHUIS, JONATHAN MIZRAHI, KALE JOHNSON, CHRISTOPHER MONROE, Joint Quantum Institute, University of Maryland Department of Physics and National Institute of Standards and Technology, College Park, Maryland 20742 — We have demonstrated ultrafast spin-motion entanglement of a single atomic ion using a short train of intense laser pulses. This pulse train gives the ion a spin-dependent kick where each spin state receives a discrete momentum kick in opposite directions. Using a series of these spin-dependent kicks we can realize a two qubit gate [1,2]. In contrast to gates using spectroscopically resolved motional sidebands, these gates may be performed faster than the trap oscillation period, making them potentially less sensitive to noise, independent of temperature, and more easily scalable to large crystals of ions. We show that multiple kicks can be strung together to create a “Schrodinger cat” like state, where the large separation between the two parts of the wavepacket allow us to accumulate the phase shift necessary for a gate in a shorter amount of time. We will present a realistic pulse scheme for a two ion gate, and our progress towards its realization.

[1] J. J. Garcí a-Ripoll, P. Zoller, and J. I. Cirac, Phys. Rev. Lett. 91, 157901 (2003).

[2] L.-M. Duan, Phys. Rev. Lett. 93, 100502 (2004).

<sup>1</sup>This work is supported by grants from the U.S. Army Research Office with funding from the DARPA OLE program, IARPA, and the MURI program; and the NSF Physics Frontier Center at JQI.

Brian Neyenhuis  
Joint Quantum Institute, University of Maryland Dept of Physics  
and National Institute of Standards and Technology,  
College Park, Maryland 20742

Date submitted: 30 Jan 2013

Electronic form version 1.4