

Abstract Submitted
for the DAMOP13 Meeting of
The American Physical Society

Progress towards loophole-free tests of the Bell inequalities by distributing ion-photon entanglement over long distances¹ LE LUO, Department of Physics, Indiana University-Purdue University Indianapolis, Indianapolis IN 46202, CHUANFENG LI, YONGJIAN HAN, YUNFENG HUANG, Key Lab of Quantum Information, CAS University of Science and Technology of China, Hefei, Anhui, China 230026 — We are developing a long term program to implement loophole-free tests of the Bell inequalities by generating entangled ion-photon pairs over long distances. Time-like interval and high detection efficiency are two essential components for a loophole-free test of the Bell inequalities. In our scheme, a pair of entangled photons (local and remote photon) from narrow-band spontaneous parametric down conversion (SPDC) will be used to produce time-like interval through a telecom fiber linking two cities separated by a distance of over 100 km in Anhui, China. Then an upconversion quantum interface will coherently transfer the qubit stored in the local SPDC photon to another photon at 370 nm which coincides the spontaneous emitted photon from trapped Yb ions. We will then implement a herald entanglement scheme by a joint measurement of both the upconversion photon and the spontaneous emitted photon which is entangled with a trapped Yb ion. Thus ion-photon entanglement can be generated between the remote SPDC photon and the trapped ion. Assuming perfect detection efficiency of the ion, a minimum detection efficiency 0.50 of the SPDC photon is required to close the detection loophole. We will present an analysis of this experimental scheme and report the current experimental progress.

¹This work is supported by the CAS (XDB01000000)

Le Luo
Department of Physics, Indiana University-Purdue
University Indianapolis, Indianapolis IN 46202

Date submitted: 30 Jan 2013

Electronic form version 1.4