

Abstract Submitted  
for the DAMOP13 Meeting of  
The American Physical Society

**Quantal determination of the mobility of ground and excited C<sup>+</sup> ions evolving in a cooled helium gas** MONCEF BOULEDROUA, Laboratoire de Physique des Rayonnements, LAMIA AISSAOUI, Physics Department - Badji Mokhtar University, KAMEL ALIOUA, Laboratoire de Physique des Rayonnements — We propose in this work to look at the mobility of C<sup>+</sup> ions moving in a neutral helium gas. The calculations are performed for the cooled buffer gas into three steps. The first step consists of calculating the interaction potentials corresponding to the dimers which dissociate into C<sup>+</sup>(<sup>2</sup>P)-He(<sup>1</sup>S) and C<sup>+</sup>(<sup>4</sup>P)-He(<sup>1</sup>S). This task is accomplished with **molpro**. Then, following the suggestions stated in a recent paper [1], we compute the energy-dependent thermophysical cross sections by using a full quantum-mechanical method, which yields in particular the quantal phase shifts. The final step aims at the use of the computed cross sections within the Viehland GRAMCHAR **fortran** code [2, 3] to get the mobility of the ions at fixed temperatures. The preliminary results are shown in the figure below.

- [1] S. Matouba, H. Tanuma, and K. Ohtsuki, J. Phys. B **41**, 145205 (2008).
- [2] L. Viehland, Chem. Phys. **179**, 71 (1994).
- [3] L. Viehland, private communications.

Moncef Bouledroua  
Laboratoire de Physique des Rayonnements

Date submitted: 22 Feb 2013

Electronic form version 1.4