

Abstract Submitted
for the DAMOP15 Meeting of
The American Physical Society

Measurement-induced generation of a macroscopic spin singlet

NAEIMEH BEHBOOD, FERRAN MARTIN CIURANA, GIORGIO COLANGELO, MARIO NAPOLITANO, ICFO-Institut de Ciències Fòniques, Mediterranean Technology Park, 08860 Castelldefels (Barcelona), Spain, GEZA TOTH¹, Department of Theoretical Physics, University of the Basque Country UPV/EHU, P.O. Box 644, E-48080 Bilbao, Spain; ROBERT J. SEWELL, ICFO-Institut de Ciències Fòniques, Mediterranean Technology Park, 08860 Castelldefels (Barcelona), Spain, MORGAN W. MITCHELL, ICFO-Institut de Ciències Fòniques, Mediterranean Technology Park, 08860 Castelldefels (Barcelona), Spain; ICREA-Institució Catalana de Recerca i — We report the production of a macroscopic spin singlet (MSS) in an atomic system [1] using collective quantum non-demolition (QND) measurement as a global entanglement generator [2]. Using an unpolarized ensemble of up to 10^6 cold atomic spins, we observe 3 dB of spin squeezing and detect entanglement with 5σ statistical significance using a generalized spin-squeezing inequality [3], indicating that at least half the atoms in the sample have formed singlets. [1] N. Behbood *et al.*, “Generation of Macroscopic Singlet States in a Cold Atomic Ensemble,” **Phys. Rev. Lett.** 113, 093601 (2014). [2] R. J. Sewell *et al.*, “Certified quantum non-demolition measurement of a macroscopic material system,” **Nat. Photon.** 7, 517 (2013). [3] G. Tóth and M.W. Mitchell, “Generation of macroscopic singlet states in atomic ensembles,” **New J. Phys.** 12, 053007 (2010).

¹IKERBASQUE, Basque Foundation for Science, E-48011 Bilbao, Spain; Wigner Research Centre for Physics, Hungarian Academy of Sciences, P.O. Box 49, H-1525 Budapest, Hungary.

Robert J. Sewell
ICFO-Institut de Ciències Fòniques, Mediterranean Technology Park,
08860 Castelldefels (Barcelona), Spain

Date submitted: 29 Jan 2015

Electronic form version 1.4