

Abstract Submitted  
for the DAMOP15 Meeting of  
The American Physical Society

**Quantum Defect Theory for Ultracold State-Resolved Chemistry<sup>1</sup>**

JISHA HAZRA, Department of Chemistry, University of Nevada, Las Vegas, NV 89154, BRANDON RUZIC, JOHN BOHN, JILA, NIST, and Department of Physics, University of Colorado, Boulder, BALAKRISHNAN NADUVALATH, Department of Chemistry, University of Nevada, Las Vegas, NV 89154 — We present a formalism for cold and ultracold atom-diatom chemical reactions that combines a quantum close-coupling method at short-range with quantum defect theory at long-range. The method yields full state-to-state rovibrationally resolved cross sections as in standard close-coupling (CC) calculations but at a considerably less computational expense. This hybrid approach exploits the simplicity of MQDT while treating the short-range interaction explicitly using quantum CC calculations. The method, demonstrated for  $D+H_2 \rightarrow HD+H$  collisions with rovibrational quantum state resolution of the HD product, is shown to be accurate for a wide range of collision energies and initial conditions. The hybrid CC-MQDT formalism may provide an alternative approach to full CC calculations for cold and ultracold reactions.

<sup>1</sup>This work was supported in part by NSF grant PHY-1205838 (N.B.) and ARO MURI grant No. W911NF-12-1-0476 (N.B. and J.L.B.)

Balakrishnan Naduvalath  
University of Nevada Las Vegas

Date submitted: 30 Jan 2015

Electronic form version 1.4