

Abstract Submitted
for the DAMOP15 Meeting of
The American Physical Society

Nuclear Spin Dependent Parity Violation in Diatomic Molecules

EMINE ALTUNTAS, JEFFREY AMMON, SIDNEY CAHN, DAVID DEMILLE, Yale University, MIKHAIL KOZLOV, Petersburg Nuclear Physics Institute, RICHARD PAOLINO, U.S. Coast Guard Academy — Nuclear spin-dependent parity violation (NSD-PV) effects arise from exchange of the Z^0 boson between electrons and the nucleus, and from interaction of electrons with the nuclear anapole moment, a parity-odd magnetic moment. The latter scales with nucleon number of the nucleus A as $A^{2/3}$, whereas the Z^0 coupling is independent of A . Thus the former is the dominant source of NSD-PV for nuclei with $A \geq 20$. We study NSD-PV effects using diatomic molecules, where signals are dramatically amplified by bringing rotational levels of opposite parity close to degeneracy in a strong magnetic field. Using a Stark-interference technique we measure the NSD-PV interaction matrix element. We present results that demonstrate statistical sensitivity to NSD-PV effects surpassing that of any previous atomic parity violation measurement, using the test system $^{138}\text{Ba}^{19}\text{F}$. We also discuss investigations of systematics due to non-reversing stray E -fields, E_{nr} together with B -field inhomogeneities, and short-term prospects for measuring the nuclear anapole moment of ^{137}Ba . In the long term, our technique is sufficiently general and sensitive to enable measurements across a broad range of nuclei.

Emine Altuntas
Yale University

Date submitted: 30 Jan 2015

Electronic form version 1.4