

Abstract Submitted
for the DAMOP15 Meeting of
The American Physical Society

Photoionization Cross Sections of P II: Theory & Measurement¹

GUILLERMO HINOJOSA, E. HERNANDEZ, A. ANTILLON, A. MORALES-MORI, A. JUAREZ, Nacional Autónoma de Mexico, A. AGUILAR, LLNL, A. COVINGTON, D. HANSTORP, K. CHATKUNCH, U of Nevada, O. GONZALEZ, U of Groninzen, D. MACALUSO, U of Montana, S. NAHAR, Ohio State U — Features of photoionization of ground and many excited states ($n \leq 10$) of P II are studied experimentally and theoretically. The ion is commonly formed in ECR ion sources and photionized. Photoionization cross sections of P II are needed for spectral analysis, but very little data are available. Experiment is done at Advanced Light Source at LLNL. Calculations are carried out in relativistic Breit-Pauli R-matrix method in close coupling approximation using a wave function of 18 levels of the core. Photoionization of the ground $3s^2 3p^2 ({}^3P_0)$ and low lying excited levels show low energy resonances with a broad feature indicating high probability of ionization. Theoretical results agree and identify the combined features of measurements. Comparison as well as other features, particularly of Seaton resonances due to photo-excitation of core, in the photoionization cross sections will be illustrated. Photoionization cross sections of 475 levels will be reported.

¹Acknowledgement of support: NSF,DOE,OSC

Sultana Nahar
Ohio State Univ - Columbus

Date submitted: 31 Jan 2015

Electronic form version 1.4