

Abstract Submitted
for the DAMOP17 Meeting of
The American Physical Society

Dark-bright soliton interactions beyond the integrable limit¹

GARYFALLIA KATSIMIGA, JAN STOCKHOFE, Center for Optical Quantum Technologies, University of Hamburg, PANAGIOTIS KEVREKIDIS, Department of Mathematics and Statistics, University of Massachusetts, Amherst, PETER SCHMELCHER, Center for Optical Quantum Technologies, University of Hamburg, THEORY GROUP OF FUNDAMENTAL PROCESSES IN QUANTUM PHYSICS TEAM — In this work we present a systematic theoretical analysis regarding dark-bright solitons and their interactions, motivated by recent advances in atomic two-component repulsively interacting Bose-Einstein condensates. In particular, we study analytically via a two-soliton ansatz adopted within a variational formulation the interaction between two dark-bright solitons in a homogeneous environment beyond the integrable regime, by considering general inter- and intra-atomic interaction coefficients. We retrieve the possibility of a fixed point in the case where the bright solitons are out of phase. As the intercomponent interaction is increased, we also identify an exponential instability of the two-soliton state, associated with a subcritical pitchfork bifurcation. The latter gives rise to an asymmetric partition of the bright soliton mass and dynamically leads to spontaneous splitting of the bound pair. In the case of the in-phase bright solitons, we explain via parsing the analytical approximations and monitoring the direct dynamics why no such pair is identified, despite its prediction by the variational analysis.

¹Deutsche Forschungsgemeinschaft in the framework of the grant SCHM 885/26-1

Garyfallia Katsimiga
Center for Optical Quantum Technologies, University of Hamburg

Date submitted: 21 Jan 2017

Electronic form version 1.4